

AGTIV®

RELIABLE INOCULANTS for CANOLA & CEREALS

- ✓ Improves photosynthesis
- ✓ Better water management
- ✓ Increases nutrient absorption



2.5 bu/ac
6.7%

Average yield increase for canola
with AGTIV® IGNITE™ L
27 sites over 5 years, Canada

3.8 bu/ac
10%

Average yield increase for cereals
with AGTIV® IGNITE™ L
8 sites over 2 years, Canada

IGNITE CROP POTENTIAL

AGTIV® is an innovative technology brand made of high-quality and proven natural active ingredients that deliver superior quality and performance for producers. Ask your local representative, or scan to learn more about the AGTIV® offer.



PROMOTED BY:



info@ptagtiv.com
PTAGTIV.COM

1 866 454-5867



LIQUID ON SEED

AGTIV® IGNITE™ L



ACTIVE INGREDIENT:

S SERENDIPITA – PTB299 Technology
Serendipita indica (formerly known as *Piriformospora indica*)
 2 x10⁶ viable spores/g

**COVERS
200
acres**

INERT INGREDIENT: Water

PARTICLE SIZE: < 1 mm (18 mesh)

Contains non-soluble particles

SIZE	COVERS	CODE
11 L (11 kg) – bag-in-box	Canola: 454 kg of seeds (1000 lb) Cereal: 9165 kg of seeds (20 205 lb)	714114

DIRECTIONS FOR USE

Ensure the seed treating equipment has been properly cleaned and calibrated and that applicator's tank is clean. Remove any filters on the treating system that are smaller than 1 mm (18 mesh) to prevent clogging. **Shake thoroughly the 11 liters bladder and add it completely to the applicator's tank.**

For canola and other Brassicaceae, one bladder of 11 liters can treat up to 454 kg (1000 lb) or 81 ha (200 acres) of seeds.

For wheat and other cereals, one bladder of 11 liters can treat up to 9165 kg (20 205 lb) or 81 ha (200 acres) of seeds. It is recommended to dilute in non-chlorinated water to reach a total volume of liquid to add between 12 to 20 ml/kg of seeds.

Spray on seeds and ensure full coverage.

BENEFITS

NUTRITIONAL ASPECTS

- (P) PHOSPHORUS – essential for energy storage and availability to cells**
 Stimulates the plant's production of phosphate transporters, enabling it to absorb phosphorus from the soil more efficiently^[1, 2].
- (N) NITROGEN – key element involved in protein and chlorophyll synthesis**
 Improves the plant's nitrogen utilization efficiency by increasing the expression of nitrate reductase, the primary enzyme used to transform absorbed mineral nitrogen into organic nitrogen, which plants use for amino acid synthesis^[3].
- (S) SULPHUR – essential for protein synthesis**
 Serendipita has high affinity sulfur transporters to absorb sulfur from the soil very efficiently, which is then transferred to the plant^[4].

ABIOTIC STRESS TOLERANCE

- Mitigates detrimental effects of water stress by stimulating the expression of protective enzymes, which play a major role in regulating reactive oxygen derivatives^[5]. These compounds accumulate under water or salt stress and can damage cell membranes and chloroplasts, affecting plant growth.
- Enhances drought tolerance by improving stomatal conductance^[6, 7].
- Improves biochemical pathways of plant partner which includes biosynthesis of prolines, organic acids and sugars, that serve as osmolytes in the cell^[6, 8, 9, 10]. This aids plants to maintain water potential gradient for the flow of water from soil into root and further to aerial parts under water deficit conditions^[11].

- [1] M. M. Aslam, J. Karanja, S.K. Bello. (2019) Piriformospora indica colonization reprograms plants to improved P-uptake, enhanced crop performance, and biotic/abiotic stress tolerance, *Physiol. Mol. Plant Pathol.* 106 (2019 April 1) 232-237
- [2] Yadav, V., Kumar, M., Deepak Kumar Deep D. et al., (2010). A phosphate transporter from the root endophytic fungus *Piriformospora indica* plays a role in phosphate transport to the host plant. *The journal of biological chemistry*, vol. 285, no. 34, p. 26535-26544
- [3] Sherameti, I., Shahollari, B., Venus, Y., Altschmied, L., Varma, A. and Oelmüller, R. (2005). The endophytic fungus *Piriformospora indica* stimulates the expression of nitrate reductase and the starch-degrading enzyme glucan-water dikinase in tobacco and Arabidopsis roots through a homeodomain transcription factor that binds to a conserved motif in their promoters. *J. Biol. Chem.* 280(28):26241-26247
- [4] Narayan, O.P., Verma, N., Jogawat, M., Dua, M. and Johri, A.K. (2021). Sulfur transfer from the endophytic fungus *Serendipita indica* improves maize growth & require the sulfate transporter SiSuIT. *Am. Soc. Of Plant Biol.* 2021. doi/10.1093/plcell/koab006/6105181.
- [5] Rong Z-Y, Jiang D-J, Cao J-L, Hashem A, Abd_Allah EF, Alsayed MF, Harsonowati W and Wu Q-S (2022) Endophytic fungus *Serendipita indica* accelerates ascorbate-glutathione cycle of white clover in response to water stress. *Front. Microbiol.* 13:967851. doi: 10.3389/fmicb.2022.967851
- [6] Jangir, P., Kanwar Shekhawat, P., Bishnoi, A, Ram, H. and Soni, P. (2021). Role of *Serendipita indica* in enhancing drought tolerance in crops. *Physiol. and Molec. Plant Pathology.* doi.org/10.1016/j.pmp.2021.101691.
- [7] Tsai, H.J., Shao, K.H., Chan, M.T., Cheng, C.P., Yeh, K.W., Oelmüller, R., and Wang, S.J. (2020). *Piriformospora indica* symbiosis improves water stress tolerance of rice through regulating stomata behavior and ROS scavenging systems. *Plant Signal. Behav.* 15(2):1722447.
- [8] Tyagi, J., Varma, A. and Pudake, R.N. (2017). Evaluation of comparative effects of arbuscular mycorrhiza (*Rhizophagus intraradices*) and endophyte (*Piriformospora indica*) association with finger millet (*Eleusine coracana*) under drought stress. *Eur. J. Soil Biol.* 81, 1-0.
- [9] Ghaffari, M.R., Mirzaei, M., Ghabouli, M., Khatibi, B. Wu, Y., Zabet-Moghaddam, M., Mohammadi-Nejad, G., Haynes, P.A., Hajirezaei, M.R., Sepehri, M. and Salekdeh, G.H. (2019). Root endophytic fungus *Piriformospora indica* improves drought stress adaptation in barley by metabolic and proteomic reprogramming. *Environ. Exp. Bot.* 157:197-210. Hosseini et al., 2017
- [10] Hosseini, F., Mosaddeghi, M.R., Dexter, A.R. and Sepehri, M. (2018). Maize water status and physiological traits as affected by root endophytic fungus *Piriformospora indica* under combined drought and mechanical stresses.
- [11] Ruiz-Lozano, J.M., Porcel, R., Azcón, C. and Aroca, R. (2012). Regulation by arbuscular mycorrhizae of the integrated physiological response to salinity in plants: new challenges in physiological and molecular studies. *J. of Exp. Bot.* 63(11):4033-4044

CANOLA

Year	Number of sites	Untreated check yield (bu/ac)	AGTIV® IGNITE™ L yield (bu/ac)	Yield increase (bu/ac)
2018	1	63.5	68.0	4.5
2019	6	44.6	47.1	2.5
2020	5	37.2	39.6	2.4
2021	8	32.5	35.0	2.5
2022	7	33.6	36.2	2.6
Total	27 sites	37.2^a	41.8^b	2.5

DURUM WHEAT

Year	Number of sites	Untreated check yield (bu/ac)	AGTIV® IGNITE™ L yield (bu/ac)	Yield increase (bu/ac)
2021	4	35.8	39.3	3.5
2022	4	40.2	44.4	4.2
Total	8 sites	38.0^a	41.8^b	3.8